

On J -orders of elements of $KO(\mathbb{C}P^m)$

Mohammad Obiedat

February 1, 2008

Abstract

Let $KO(\mathbb{C}P^m)$ be the KO -ring of the complex projective space $\mathbb{C}P^m$. By means of methods of rational D -series [4], a formula for the J -orders of elements of $KO(\mathbb{C}P^m)$ is given. Explicit formulas are given for computing the J -orders of the canonical generators of $KO(\mathbb{C}P^m)$ and the J -order of any complex line bundle over $\mathbb{C}P^m$.

1 Introduction

Let X be a connected finite-dimensional CW complex. Let $KO(X)$ be the KO -ring of X and $\tilde{K}O(X)$ (resp. $\tilde{K}SO(X)$) be the subgroup of $KO(X)$ of elements (resp. orientable elements) of virtual dimension zero. For a real vector bundle E over X , let $S(E)$ be the sphere bundle associated to E with respect to some inner product on E . Let $JO(X) = KO(X)/TO(X)$ be the J -group of X , where $TO(X) = \{E - F \in \tilde{K}O(X) : S(E \oplus n) \text{ is fibre homotopy equivalent to } S(F \oplus n) \text{ for some } n \in \mathbb{N}\}$. Then by Adams [1] and Quillen [9], it is shown that

$$TO(X) = \{ x \in \tilde{K}SO(X) : \text{there exists } u \in \tilde{K}SO(X) \text{ such that} \\ \theta_k(x) = \frac{\psi^k(1+u)}{1+u} \text{ in } 1 + \tilde{K}SO(X) \otimes \mathbb{Q}_k \text{ for all } k \in \mathbb{N} \} \quad (1)$$

where $\theta_k : \tilde{K}SO(X) \rightarrow 1 + \tilde{K}SO(X) \otimes \mathbb{Q}_k$ is the Bott exponential map, ψ^k is the Adams operation, $\mathbb{Q}_k = \{n/k^m : n, m \in \mathbb{Z}\}$, and $1 + \tilde{K}SO(X) \otimes \mathbb{Q}_k$ is the multiplicative group of elements $1 + w$ with $w \in \tilde{K}SO(X) \otimes \mathbb{Q}_k$.

Now, X is connected implies that $KO(X) = \tilde{K}O(X) \oplus \mathbb{Z}$. So, $JO(X) = \tilde{J}O(X) \oplus \mathbb{Z}$ where $\tilde{J}O(X) = \tilde{K}O(X)/TO(X)$. For $x \in KO(X)$, the J -order of x is the order of $x + TO(X)$ in $JO(X)$. By Atiyah [3], $\tilde{J}O(X)$ is a finite group. Hence, $x \in KO(X)$ has a finite J -order if and only if $x \in \tilde{K}O(X)$. Let $y_m = r\xi_m(\mathbb{C}) - 2$ where $\xi_m(\mathbb{C})$ is the complex Hopf line bundle over the complex projective space $\mathbb{C}P^m$. Every element of $\tilde{K}O(\mathbb{C}P^m)$ has the form

$P_m(y_m; m_1, \dots, m_s) = m_1 y_m + \dots + m_s y_m^s$ for some $m_i \in \mathbb{Z}$ and $s \in \mathbb{N}$. The purpose of this paper is to compute the J -order of $P_m(y_m; m_1, \dots, m_s)$ which we denote by $b_m(P_m(y_m; m_1, \dots, m_s))$. In addition to their self importance, these orders are of great importance in geometric topology, for instance, it is well-known that the Stiefel fibration $U(n)/U(n-m-1) \longrightarrow S^{2n-1}$ has a cross-section if and only if n is a multiple of $b_m(P_m(y_m; 1, 0, \dots, 0))$, computed by Adams-Walker [2]. Dibağ [5] has used (1) to give another proof of Lam's results [6], we rediscover his proof as a special case of Example 2 below.

In section 2, we first obtain a useful formula for $\theta_p(y_m^n)$ for $n = 1, \dots, s$. Then we use (1) and some facts of rational D -series [4] to give, in Theorem 2.5, a formula for $b_m(P_m(y_m; m_1, \dots, m_s))$. The formula given in Theorem 2.5 involves many difficulties and one can obtain a little information about the range of the J -orders of elements of $\tilde{K}O(\mathbb{C}P^m)$. So instead, we use a well-known computations of the J -order of y_m to obtain, in Theorem 2.6, upper and lower bounds for $b_m(P_m(y_m; m_1, \dots, m_s))$.

There are two important examples in mind, namely the J -orders of the canonical generators of $\tilde{K}O(\mathbb{C}P^m)$ and the J -orders of complex line bundles over $\mathbb{C}P^m$. In section 3 we first obtain an explicit formula for the J -orders of y_m^n for $n = 2, 3$, or 4 , we conjecture that the formula is liable to be true for $k = 5, \dots, t$. Then, we give an explicit formula for the J -order of any complex line bundle over $\mathbb{C}P^m$.

2 The J -order of $P_m(y_m; m_1, \dots, m_s) \in \tilde{K}O(\mathbb{C}P^m)$

Let L be a non-trivial complex line bundle over $\mathbb{C}P^m$. Then $L \cong \xi_m(\mathbb{C})^n$ or $L \cong \overline{\xi_m(\mathbb{C})}^n$ for some $n \in \mathbb{N}$, where $\overline{\xi_m(\mathbb{C})}$ denotes the conjugate bundle to $\xi_m(\mathbb{C})$. Let $KU(\mathbb{C}P^m)$ be the KU -ring of $\mathbb{C}P^m$. In Lemma 2.1, we find the image of $\xi_m(\mathbb{C})^n$ under the realification homomorphism

$$r : KU(\mathbb{C}P^m) \rightarrow KO(\mathbb{C}P^m).$$

Note that $r(\overline{\xi_m(\mathbb{C})}^n) = r(\xi_m(\mathbb{C})^n)$.

Recall that from [2], $KO(\mathbb{C}P^m)$ is a truncated polynomial ring over the integers generated by y_m with the following relations:

$$\begin{aligned} y_m^{t+1} &= 0 \text{ if } m = 2t; \\ 2y_m^{2t+1} &= 0, \quad y_m^{2t+2} = 0 \text{ if } m = 4t + 1; \\ y_m^{2t+2} &= 0 \text{ if } m = 4t + 3. \end{aligned}$$

For each $m \in \mathbb{N}$, let

$$d_m = \begin{cases} t & \text{if } m = 2t \\ 2t + 1 & \text{if } m = 4t + 1 \text{ or } m = 4t + 3. \end{cases}$$

For each $r, s \in \mathbb{N}$ with $r \geq s$, let $b_{r,s}$ be the coefficient of $(\xi_m(\mathbb{C})^s + \overline{\xi_m(\mathbb{C})})^s$ in $(\xi_m(\mathbb{C}) + \overline{\xi_m(\mathbb{C})} - 2)^r$. Using the fact that $\xi_m(\mathbb{C}) \overline{\xi_m(\mathbb{C})} = 1$, we easily see that:

$$b_{r,s} = (-2)^{r-s} \binom{r}{s} + \sum_{\substack{j=s+1 \\ j=s+2s_j}}^r (-2)^{r-j} \binom{r}{j} \binom{j}{s_j}. \quad (2)$$

Further, $b_{r,0}$ is the constant term of $(\xi_m(\mathbb{C}) + \overline{\xi_m(\mathbb{C})} - 2)^r$.

Let $n \in \mathbb{N}$. For each $s = 1, \dots, n$, define $d_{n,s}$ by the recurrence relation $d_{n,n} = 1$ and for $s = n-1, n-2, \dots, 1$

$$d_{n,s} = -(d_{n,s+1}b_{s+1,s} + d_{n,s+2}b_{s+2,s} + \dots + d_{n,n}b_{n,s}). \quad (3)$$

Convention. Let $f \in \mathbb{Z}[[y_m]]$, where $\mathbb{Z}[[y_m]]$ is the ring of formal power series with coefficients in $\mathbb{Z}$. If we consider f as an element of $KO(\mathbb{C}P^m)$, then we implicitly mean that $f \pmod{y_m^{d_m+1}}$.

Lemma 2.1 *Let $n, m \in \mathbb{N}$. Then*

$$(i) \quad r(\xi_m(\mathbb{C})^n) = d_{n,1}y_m + d_{n,2}y_m^2 + \dots + d_{n,n}y_m^n + 2.$$

$$(ii) \quad \psi^n(y_m) = d_{n,1}y_m + d_{n,2}y_m^2 + \dots + d_{n,n}y_m^n, \text{ and for } k = 2, \dots, d_m,$$

$$\psi^n(y_m^k) = d_{kn,1}y_m + \dots + d_{kn,kn}y_m^{kn} - d_{k,1}\psi^n(y_m) - \dots - d_{k,k-1}\psi^n(y_m^{k-1}).$$

$$(iii) \quad d_{n,1} = n^2.$$

Proof. (i) Let $c : KO(\mathbb{C}P^m) \rightarrow KU(\mathbb{C}P^m)$ be the complexification homomorphism. $c r(\xi_m(\mathbb{C})^n) = \xi_m(\mathbb{C})^n + \overline{\xi_m(\mathbb{C})}^n$. On the other hand, by (2),(3) and the fact that $y_m^n = 0$ for $n > d_m$, we have

$$c(d_{n,1}y_m + \dots + d_{n,n}y_m^n + 2) = \xi_m(\mathbb{C})^n + \overline{\xi_m(\mathbb{C})}^n.$$

Using the fact that c is a monomorphism for $m = 2t$ and $m = 4t + 3$, we get

$$r(\xi_m(\mathbb{C})^n) = d_{n,1}y_m + \dots + d_{n,n}y_m^n + 2.$$

To prove the case $m = 4t + 1$, let $i : \mathbb{C}P^{4t+1} \rightarrow \mathbb{C}P^{4t+2}$ be the inclusion map. Then $i^* : KO(\mathbb{C}P^{4t+2}) \rightarrow KO(\mathbb{C}P^{4t+1})$ is an epimorphism and maps $r(\xi_{4t+2}(\mathbb{C})^n)$ to $r(\xi_{4t+1}(\mathbb{C})^n)$. Hence,

$$\begin{aligned} r(\xi_{4t+1}(\mathbb{C})^n) &= i^*(r(\xi_{4t+2}(\mathbb{C})^n)) = i^*(d_{n,1}y_{4t+2} + \dots + d_{n,n}y_{4t+2}^n + 2) \\ &= d_{n,1}y_{4t+1} + d_{n,2}y_{4t+1}^2 + \dots + d_{n,n}y_{4t+1}^n + 2. \end{aligned}$$

(ii) Let $l \in \{1, \dots, d_m\}$. By (i),

$$\begin{aligned} d_{ln,1}y_m + \dots + d_{ln,ln}y_m^{ln} &= r(\xi_m(\mathbb{C})^{ln}) - 2 = r\psi^n(\xi_m(\mathbb{C})^l) - 2 \\ &= \psi^n(r\xi_m(\mathbb{C})^l) - 2 = \psi^n(d_{l,1}y_m + \dots + d_{l,l}y_m^l). \end{aligned}$$

The result follows.

(iv) $d_{n,1}$ is the constant term of

$$\begin{aligned} \frac{cr(\xi_m(\mathbb{C})^n) - 2}{\xi_m(\mathbb{C}) + \overline{\xi_m(\mathbb{C})} - 2} &= \frac{\xi_m(\mathbb{C})^n + \overline{\xi_m(\mathbb{C})}^n - 2}{\xi_m(\mathbb{C}) + \overline{\xi_m(\mathbb{C})} - 2} = \\ &(\xi_m(\mathbb{C})^{n-1} + \xi_m(\mathbb{C})^{n-2} + \dots + \xi_m(\mathbb{C}) + 1)((\overline{\xi_m(\mathbb{C})}^{n-1} + \overline{\xi_m(\mathbb{C})}^{n-2} + \dots + \overline{\xi_m(\mathbb{C})} + 1)). \end{aligned}$$

Hence,

$$\begin{aligned} d_{n,1} &= n + 2(n-1) + 2(n-2) + \dots + 2 \\ &= n + 2((n-1) + (n-2) + \dots + 1) = n + 2\left(\frac{n(n-1)}{2}\right) = n^2. \end{aligned}$$

This completes the proof of Lemma 2.1.

Now, we use the above lemma to find $\theta_p(y_m^n) \in 1 + \tilde{K}O(\mathbb{C}P^m) \otimes \mathbb{Q}_p$. For each $n, m, p \in \mathbb{N}$ with $n \leq d_m$, let

$$A(p; n, m) = \left(\frac{\psi^p(d_{n,1} + \dots + d_{n,n}y_m^{n-1})}{d_{n,1} + \dots + d_{n,n}y_m^{n-1}} \right)^{\frac{1}{2}} \in 1 + \tilde{K}O(\mathbb{C}P^m) \otimes \mathbb{Q}_p,$$

and for $n \geq 2$, let

$$B(p; n, m) = (\theta_p(y_m)^{d_{n,1}-1} \theta_p(y_m^2)^{d_{n,2}} \dots \theta_p(y_m^{n-1})^{d_{n,n-1}})^{-1} \in 1 + \tilde{K}O(\mathbb{C}P^m) \otimes \mathbb{Q}_p.$$

Theorem 2.2 *Let $p \geq 2$ and $m = 2t$ for some $t \geq 1$. Then*

$$(i) \quad \theta_p(y_m) = \left(\frac{\psi^p(y_m)}{p^2 y_m} \right)^{\frac{1}{2}}.$$

$$(ii) \quad \theta_p(y_m^n) = A(p; n, m)B(p; n, m), \text{ for each } 2 \leq n \leq t.$$

Proof. (i) This is Lemma 5.4 of [7].

(ii) If η is a complex $4n$ -dimensional vector bundle over a finite CW complex X such that $\wedge^{4n}\eta = 1$, then $c\theta_p(r\eta) = \theta_p(\eta)$. Let $\eta = 2\xi_m(\mathbb{C})^n + 2\overline{\xi_m(\mathbb{C})}^n$. Then η is a 4 -dimensional complex vector bundle over $\mathbb{C}P^m$ with

$$\wedge^4(\eta) = \wedge^2(2\xi_m(\mathbb{C})^n) \wedge^2(2\overline{\xi_m(\mathbb{C})}^n) = 1.$$

Hence, $c\theta_p(r\eta) = \theta_p(\eta)$. By Lemma 2.1,

$$\xi_m(\mathbb{C})^n + \overline{\xi_m(\mathbb{C})}^n - 2 = cr(\xi_m(\mathbb{C})^n) - 2 = c(d_{n,1}y_m + \dots + d_{n,n}y_m^n).$$

Also,

$$\begin{aligned} r\eta &= 2r(cr(\xi_m(\mathbb{C})^n)) = 2rc(r(\xi_m(\mathbb{C})^n)) \\ &= 4r(\xi_m(\mathbb{C})^n) = 4d_{n,1}y_m + \cdots + 4d_{n,n}y_m^n + 8. \end{aligned}$$

Thus,

$$c\theta_p(r\eta) = c((p\theta_p(y_m)^{d_{n,1}}\theta_p(y_m^2)^{d_{n,2}} \cdots \theta_p(y_m^{n-1})^{d_{n,n-1}}\theta_p(y_m^n)^4).$$

On the other hand,

$$\begin{aligned} \theta_p(\eta) &= \left(\frac{\xi_m(\mathbb{C})^{np} + \overline{\xi_m(\mathbb{C})}^{np} - 2}{\xi_m(\mathbb{C})^n + \overline{\xi_m(\mathbb{C})}^n - 2} \right)^2 = \left(\frac{\psi^p(\xi_m(\mathbb{C})^n + \overline{\xi_m(\mathbb{C})}^n - 2)}{\xi_m(\mathbb{C})^n + \overline{\xi_m(\mathbb{C})}^n - 2} \right)^2 \\ &= \left(\frac{c(\psi^p(y_m))c(\psi^p(d_{n,1} + d_{n,2}y_m + \cdots + d_{n,n}y_m^{n-1}))}{c(y_m)c(d_{n,1} + d_{n,2}y_m + \cdots + d_{n,n}y_m^{n-1})} \right)^2. \end{aligned}$$

By (i),

$$c\left(\frac{\psi^p(y_m)}{y_m}\right) = c(p^2\theta_p(y_m)^2).$$

Hence,

$$\theta_p(\eta) = c\left((p\theta_p(y_m))^4\left(\frac{\psi^p(d_{n,1} + d_{n,2}y_m + \cdots + d_{n,n}y_m^{n-1})}{d_{n,1} + d_{n,2}y_m + \cdots + d_{n,n}y_m^{n-1}}\right)^2\right).$$

Now, c is a monomorphism implies that

$$\begin{aligned} &p\theta_p(y_m)^{d_{n,1}}\theta_p(y_m^2)^{d_{n,2}} \cdots \theta_p(y_m^{n-1})^{d_{n,n-1}}\theta_p(y_m^n) \\ &= p\theta_p(y_m) \left(\frac{\psi^p(d_{n,1} + d_{n,2}y_m + \cdots + d_{n,n}y_m^{n-1})}{d_{n,1} + d_{n,2}y_m + \cdots + d_{n,n}y_m^{n-1}} \right)^{\frac{1}{2}}. \end{aligned}$$

Hence,

$$\theta_p(y_m^n)B(p; n, m)^{-1} = A(p; n, m).$$

This completes the proof of Theorem 2.2.

REMARK. By Using a method similar to that used in proving Lemma 2.1 when $m = 4t + 1$, we easily obtain a similar formula for $\theta_p(y_m^n)$ when m is an odd integer.

Corollary 2.3 $\theta_p \circ \psi^n = \psi^n \circ \theta_p$ on $\widetilde{KO}(\mathbb{CP}^m)$ for all $m, n, p \in \mathbb{N}$.

Proof. Clearly, we only need to show that $\theta_p \circ \psi^n(y_m^k) = \psi^n \circ \theta_p(y_m^k)$ for each $k = 1, \dots, d_m$. By induction on k , if $k = 1$ then by Theorem 2.2 and Lemma 2.1 (ii),

$$\theta_p(\psi^n(y_m)) = \theta_p(d_{n,1}y_m + \cdots + d_{n,n}y_m^n)$$

$$\begin{aligned}
&= \left(\frac{\psi^p(d_{n,1} + d_{n,2}y_m \cdots + d_{n,n}y_m^{n-1})}{d_{n,1} + d_{n,2}y_m + \cdots + d_{n,n}y_m^{n-1}} \right)^{\frac{1}{2}} \theta_p(y_m) \\
&= \left(\frac{\psi^p(\psi^n(y_m))}{p^2\psi^n(y_m)} \right)^{\frac{1}{2}} = \psi^n \left(\frac{\psi^p(y_m)}{p^2y_m} \right)^{\frac{1}{2}} = \psi^n(\theta_p(y_m)).
\end{aligned}$$

Now, suppose $\theta_p(\psi^n(y_m^l)) = \psi^n(\theta_p(y_m^l))$ for $l = 1, \dots, k-1$. Then

$$\begin{aligned}
\theta_p(\psi^n(y_m^k)) &= \theta_p(d_{kn,1}y_m + \cdots + d_{kn,kn}y_m^{kn} - d_{k,1}\psi^n(y_m) - \cdots - d_{k,k-1}\psi^n(y_m^{k-1})) \\
&= \theta_p(y_m)^{d_{kn,1}} \cdots \theta_p(y_m^{kn})^{d_{kn,kn}} \theta_p(\psi^n(y_m))^{-d_{k,1}} \cdots \theta_p(\psi^n(y_m^{k-1}))^{-d_{k,k-1}}.
\end{aligned}$$

On the other hand, $\psi^n(\theta_p(y_m^k)) = \psi^n(A(p; k, m)B(p; k, m))$. So, we only need to show that

$$\begin{aligned}
&\theta_p(y_m)^{d_{kn,1}} \cdots \theta_p(y_m^{kn})^{d_{kn,kn}} = A(p; k, m)\psi^n(\theta_p(y_m)). \\
&\theta_p(y_m)^{d_{kn,1}} \cdots \theta_p(y_m^{kn})^{d_{kn,kn}} = \left(\frac{\psi^p(d_{kn,1} + d_{kn,2}y_m^2 \cdots + d_{kn,kn}y_m^{kn-1})}{d_{kn,1} + d_{kn,2}y_m + \cdots + d_{kn,kn}y_m^{kn-1}} \right)^{\frac{1}{2}} \theta_p(y_m) \\
&= \left(\frac{\psi^p(d_{kn,1}y_m + d_{kn,2}y_m^2 \cdots + d_{kn,kn}y_m^{kn})}{p^2(d_{kn,1}y_m + d_{kn,2}y_m^2 + \cdots + d_{kn,kn}y_m^{kn})} \right)^{\frac{1}{2}} = \left(\frac{\psi^p(\psi^{kn}(y_m))}{p^2\psi^{kn}(y_m)} \right)^{\frac{1}{2}} \\
&= \left(\psi^n \left(\frac{\psi^p(\psi^k(y_m))}{p^2\psi^k(y_m)} \right) \right)^{\frac{1}{2}} = A(p; k, m)\psi^n(\theta_p(y_m)).
\end{aligned}$$

The result follows.

Let $s = d_m$ and $P_m(y_m; m_1, \dots, m_s) = m_1y_m + \cdots + m_sy_m^s \in \tilde{K}O(\mathbb{C}P^m)$. Let $b_m(P_m(y_m; m_1, \dots, m_s))$ be the J -order of $P_m(y_m; m_1, \dots, m_s)$, that is the order of $P_m(y_m; m_1, \dots, m_s) + TO(\mathbb{C}P^m)$ in $\tilde{J}O(\mathbb{C}P^m)$. By using the same method of Proposition 5.7 of [7], we easily see that

$$b_{4t+3}(P_{4t+3}(y_{4t+3}; m_1, \dots, m_{2t+1})) = b_{4t+2}(P_{4t+2}(y_{4t+2}; m_1, \dots, m_{2t+1})),$$

and $b_{4t+1}(P_{4t+1}(y_{4t+1}; m_1, \dots, m_{2t+1}))$

$$= \begin{cases} b_{4t}(P_{4t}(y_{4t}; m_1, \dots, m_{2t})) & \text{if } m_{2t+1} = 0 \\ lcm\{b_{4t}(P_{4t}(y_{4t}; m_1, \dots, m_{2t})), 2\} & \text{if } m_{2t+1} = 1. \end{cases}$$

Therefore, in the remainder of this paper we shall assume that $m = 2t$ for some $t \geq 1$, unless otherwise indicated.

To compute $b_m(P_m(y_m; m_1, \dots, m_t))$, we use the notion of rational D -series introduced and developed in [4]. Let $\mathbb{Q}[[x]]$ be the ring of formal power series with coefficients in $\mathbb{Q}$ and $\mathbb{Q}^*[[x]] = \{f(x) \in \mathbb{Q}[[x]] : f(0) = \pm 1\}$. Let

$$f(x) = \pm 1 + \sum_{i \geq 1} a_i x^i \in \mathbb{Q}^*[[x]].$$

For each $k \geq 1$, let $e_k(f)$ denote the smallest positive integer e_k such that $(f(x))^{e_k} \in \mathbb{Z}[[x]](\text{mod } x^{k+1})$. Let $S_k(f)$ be the set of all primes dividing the denominators of the coefficients a_i for $i = 1, \dots, k$. For a rational number q , let $\nu_p(q)$ be the exponent of p in the prime factorization of q and let $D(q)$ be the denominator of q in its lowest term. For convenience, we assume that $\nu_p(0) = -\infty$ and $D(0) = 1$. It follows from Lemma 1.3 [4] that $p \in S_k(f)$ if and only if $\nu_p(e_k(f)) > 0$.

For each prime p , let $\alpha_p, \beta_p \in \mathbb{Z}^+$, and let $\alpha = (\alpha_2, \alpha_3, \alpha_5, \dots)$, and $\beta = (\beta_2, \beta_3, \beta_5, \dots)$. A series $f = \pm 1 + \sum_{i \geq 1} a_i x^i \in \mathbb{Q}^*[[x]]$ is called a rational D -series of type (α, β) if $\nu_p(a_{\alpha_p}) = -\beta_p$ and $\nu_p(a_k) \geq -\beta_p[\frac{k}{\alpha_p}]$ for each $k > \alpha_p$ with $a_k \neq 0$. A rational D -series f is called strict at a prime p if $\nu_p(a_k) > -\beta_p[\frac{k}{\alpha_p}]$ for each $k > \alpha_p$ with $a_k \neq 0$. If f is strict at p then, by Theorem 3.5 of [4],

$$\nu_p(e_k(f)) = \max\{\beta_p r + \nu_p(r) : 0 \leq r \leq [\frac{k}{\alpha_p}]\}.$$

With these facts on hand, we return to our problem.

Let $\mathbb{Z}_{(p)} = \{\frac{r}{s} : r, s \in \mathbb{Z} \text{ with } \nu_p(s) = 0\}$ be the localization of $\mathbb{Z}$ at p . The following lemma is (5.2) and Lemma 5.5 of [7] with minor changes.

Lemma 2.4 (i) *Let $1 + u$ be an element of $\mathbb{Q}^*[[y_m]](\text{mod } y_m^{t+1})$. Then*

$$\frac{\psi^p(1 + u)}{1 + u} \in \mathbb{Z}_{(p)}^*[[y_m]](\text{mod } y_m^{t+1})$$

if and only if $u \in \mathbb{Z}_{(p)}[[y_m]](\text{mod } y_m^{t+1})$ with $u(0) = 0$.

(ii) $(\theta_p(y_m))^h \in \mathbb{Z}_{(p)}^*[[y_m]](\text{mod } y_m^{t+1})$ if and only if

$$\nu_p(h) \geq \nu_p(b_m(y_m)) = \max\{s + \nu_p(s) : 0 \leq s \leq [\frac{m}{p-1}]\}.$$

Now, we compute $b_m(P_m(y_m; m_1, \dots, m_t))$. For each $n = 1, \dots, t$, let $\theta_p(y_m^n) = 1 + \alpha_{n,1}y_m + \dots + \alpha_{n,t}y_m^t$ where $\alpha_{n,i}$ is the coefficient of y_m^i given by Theorem 2.2. According to (1) $b_m(P_m(y_m; m_1, \dots, m_t))$ is the smallest positive integer h such that

$$\theta_p(hP_m(y_m; m_1, \dots, m_t)) = \frac{\psi^p(1 + u)}{1 + u} \text{ in } \tilde{KO}(\mathbb{CP}^m) \otimes \mathbb{Q}$$

for some $u \in \tilde{KO}(\mathbb{CP}^m)$ and all primes p . Let

$$\begin{aligned} \beta_m(y_m; m_1, \dots, m_t) &= \theta_p(P_m(y_m; m_1, \dots, m_t)) = \theta_p(y_m)^{m_1} \dots \theta_p(y_m^t)^{m_t} \\ &= 1 + \alpha_1(m_1, \dots, m_t)y_m + \dots + \alpha_t(m_1, \dots, m_t)y_m^t, \end{aligned}$$

for some $\alpha_i(m_1, \dots, m_t) \in \mathbb{Q}$. Then

$$\theta_p(hP_m(y_m; m_1, \dots, m_t)) = \beta_m(y_m; m_1, \dots, m_t)^h.$$

Let

$$\beta_m(y_m; m_1, \dots, m_t)^h = \frac{\psi^p(1+u)}{1+u}$$

for some $u \in \tilde{KO}(\mathbb{C}P^m)$. Then $\beta_m(y_m; m_1, \dots, m_t)^h$ has integer coefficients. Hence

$$\nu_p(h) \geq \nu_p(e_t(\beta_m(y_m; m_1, \dots, m_t))).$$

On the other hand, if $b = e_t(\beta_m(y_m; m_1, \dots, m_t))$ then

$$\beta_m(y_m; m_1, \dots, m_t)^b = \left(\frac{\psi^p(1+u)}{1+u}\right)^{\frac{b}{h}} = \frac{\psi^p(1+w)}{1+w}$$

for some $w \in \mathbb{Q}^*[y_m](\text{mod } y_m^{t+1})$. $\beta_m(y_m; m_1, \dots, m_t)^b \in \mathbb{Z}^*[y_m](\text{mod } y_m^{t+1})$ implies that

$$\frac{\psi^p(1+w)}{1+w} \in \mathbb{Z}^*[y_m](\text{mod } y_m^{t+1}).$$

So, by Lemma 2.4 $w \in \tilde{KO}(\mathbb{C}P^m)$ and hence $\nu_p(h) \leq \nu_p(b)$. By Corollary 1.3 of [4], $e_t(\beta_m(y_m; m_1, \dots, m_t)) = D(b_1) \cdots D(b_t)$ where $b_1 = \alpha_1(m_1, \dots, m_t)$ and for $k = 1, \dots, t$, b_k is the coefficient of y_m^k in $\beta_m(y_m; m_1, \dots, m_t)^{D(b_1) \cdots D(b_{k-1})}$. So far, we have proved:

Theorem 2.5 $b_m(P_m(y_m; m_1, \dots, m_t)) = D(b_1) \cdots D(b_t)$.

Although Theorem 2.5 gives $b_m(P_m(y_m; m_1, \dots, m_t))$ by a formula, it is difficult to use this formula to find $b_m(P_m(y_m; m_1, \dots, m_t))$ for specific values of $m_1, \dots, m_t$, because one needs first to find the coefficients of y_m^k in $\theta_p(y_m)^{m_1} \cdots \theta_p(y_m^t)^{m_t}$ and then to find $e_t(\beta_m(y_m; m_1, \dots, m_t))$ which involves tedious calculations. So, alternatively, we next try to obtain information about $b_m(P_m(y_m; m_1, \dots, m_t))$ by using what we know about $b_m(y_m)$.

By Theorem 2.2, we directly obtain

$$\theta_p(y_m^k) = \frac{\alpha_k}{\theta_p(y_m)^{N_k}} \text{ where } \alpha_2 = \left(\frac{\psi^p(d_{2,1} + y_m)}{d_{2,1} + y_m}\right)^{\frac{1}{2}}, N_2 = d_{2,1} - 1$$

and for $k = 3, \dots, t$,

$$\alpha_k = \left(\frac{\psi^p(d_{k,1} + d_{k,2}y_m + \cdots + d_{k,k}y_m^{k-1})}{d_{k,1} + d_{k,2}y_m + \cdots + d_{k,k}y_m^{k-1}}\right)^{\frac{1}{2}} \alpha_2^{-d_{k,2}} \alpha_3^{-d_{k,3}} \cdots \alpha_{k-1}^{-d_{k,k-1}},$$

$$N_k = (d_{k,1} - 1) - N_2 d_{k,2} - \cdots - N_{k-1} d_{k,k-1}. \quad (4)$$

For each $m_1, \dots, m_t \in \mathbb{Z}$, let

$$E(m_1, \dots, m_t) = \text{lcm}\{e_t(\alpha_2^{m_2}), \dots, e_t(\alpha_t^{m_t})\},$$

$$N(m_1, \dots, m_t) = m_1 - m_2 N_2 - \dots - m_t N_t,$$

$$L(p; m_1, \dots, m_t) = \nu_p(b_m(y_m)) - \nu_p(N(m_1, \dots, m_t)) - \nu_p(E(m_1, \dots, m_t)),$$

and

$$U(p; m_1, \dots, m_t) = \max\{\nu_p(b_m(y_m)) - \nu_p(N(m_1, \dots, m_t)), \nu_p(E(m_1, \dots, m_t))\}.$$

Theorem 2.6 *Let $P_m(y_m; m_1, \dots, m_t) = m_1 y_m + \dots + m_t y_m^t \in \tilde{K}O(\mathbb{C}P^m)$. Then*

$$L(p; m_1, \dots, m_t) \leq \nu_p(b_m(P_m(y_m; m_1, \dots, m_t))) \leq U(p; m_1, \dots, m_t).$$

Proof. Let $h = b_m(P_m(y_m; m_1, \dots, m_t))$. Then

$$\theta_p(m_1 y_m + \dots + m_t y_m^t)^h = \frac{\psi^p(1+u)}{1+u}$$

for some $u \in \tilde{K}O(\mathbb{C}P^m)$ and all primes p . So,

$$\theta_p(y_m)^{N(m_1, \dots, m_t)h} = \frac{\psi^p(1+u)}{1+u} \alpha_2^{-m_2 h} \dots \alpha_t^{-m_t h}.$$

Thus,

$$\theta_p(y_m)^{N(m_1, \dots, m_t)E(m_1, \dots, m_t)h}$$

has integer coefficients. Hence, by Lemma 2.4 (ii)

$$\nu_p(N(m_1, \dots, m_t)) + \nu_p(E(m_1, \dots, m_t)) + \nu_p(h) \geq \nu_p(b_m(y_m)),$$

namely $\nu_p(h) \geq L(p; m_1, \dots, m_t)$. On the other hand, let $b \in \mathbb{N}$ such that $\nu_p(b) = U(p; m_1, \dots, m_t)$. Then

$$\theta_p(P_m(y_m; m_1, \dots, m_t))^b = \left(\frac{\psi^p(1+u)}{1+u} \right)^{\frac{b}{h}}.$$

So,

$$\theta_p(y_m)^{N(m_1, \dots, m_t)b} = \left(\frac{\psi^p(1+u)}{1+u} \right)^{\frac{b}{h}} \alpha_2^{-m_2 b} \dots \alpha_t^{-m_t b}.$$

Let

$$\left(\frac{\psi^p(1+u)}{1+u} \right)^{\frac{b}{h}} = \frac{\psi^p(1+w)}{1+w}$$

for some $w \in \mathbb{Q}^*[[y_m]](\text{mod } y_m^{t+1})$ with $w(0) = 0$. Now,

$$\alpha_2^{-m_2 b} \dots \alpha_t^{-m_t b} \text{ and } \theta_p(y_m)^{N(m_1, \dots, m_t)b}$$

have integer coefficients. Hence,

$$\frac{\psi^p(1+w)}{1+w}$$

has integer coefficients, which implies that $w \in \mathbb{Z}^*[[y_m]](\text{mod } y_m^{t+1})$. Hence,

$$\nu_p(h) \leq \nu_p(b).$$

This completes the proof of Theorem 2.6.

Corollary 2.7 *Let $P_m(y_m; m_1, \dots, m_t) = m_1 y_m + \dots + m_t y_m^t \in \tilde{K}O(\mathbb{C}P^m)$. Let s be the smallest positive integer such that $m_i = 0$ for all $i > s$, if $m_t \neq 0$, let $s = t$. Then*

$$\nu_p(b_m(P_m(y_m; m_1, \dots, m_t))) = \max\{\nu_p(b_m(y_m)) - \nu_p(N(m_1, \dots, m_t)), 0\}$$

for all $p > s$.

Proof. Using Lemma 2.1 (iii), we easily see that $S_t(\alpha_k^{m_k}) \subseteq \{2, 3, \dots, k\}$ for each $k = 2, \dots, t$. So, if $p > s$ then $\nu_p(E(m_1, \dots, m_t)) = 0$. Now, the result follows from Theorem 2.6.

3 Two Important Examples

Let $m = 2t$. Then

$$\tilde{JO}(\mathbb{C}P^m) = \langle \alpha_1 = y_m + TO(\mathbb{C}P^m), \dots, \alpha_t = y_m^t + TO(\mathbb{C}P^m) \rangle.$$

In Example 1, we give a simple formula for the J -orders of α_2, α_3 , and α_4 . We conjecture that our formula is liable to be true for the J -orders of α_k , $k = 5, \dots, t$.

Let $L \cong \xi_m(\mathbb{C})^n$ for some $n \in \mathbb{N}$. By the J -order of L we mean the order of $r\xi_m(\mathbb{C})^n - 2 + TO(\mathbb{C}P^m)$ in $JO(\mathbb{C}P^m)$. Lam [6] has used complex K-theory to find the J -order of L when n is a prime power. Also, Dibağ [5] has used (1) to give another proof of Lam's results. In Example 2, we give a simple formula for the J -order of L for each $n \in \mathbb{N}$.

EXAMPLE 1. For each $k = 2, \dots, t$, the J -order of $\alpha_k = y_m^k + TO(\mathbb{C}P^m)$ is

$$b_m(P_m(y_m; 0, \dots, 0, m_k = 1, 0, \dots, 0)).$$

So, by Corollary 2.7, if $p > k$ then

$$\nu_p(b_m(y_m^k)) = \nu_p(b_m(y_m)) - \nu_p(N_k).$$

The first few values of N_k are $N_2 = 3$, $N_3 = -2.5$, $N_4 = 5.7$, $N_5 = -2.3^2.7$, $N_6 = 2.3.7.11$. By inspection, if $p > k$ then $\nu_p(N_k) = \lfloor \frac{2(k-1)}{p-1} \rfloor$ for each $k = 2, \dots, 6$.

In [8], we proved that if $p = 2$, or 3 , then

$$\nu_p(b_m(y_m^k)) = \max\{r - \lfloor \frac{2(k-1)}{p-1} \rfloor + \nu_p(r) : \lfloor \frac{2k}{p-1} \rfloor \leq r \leq \lfloor \frac{m}{p-1} \rfloor\}$$

for each $k = 1, \dots, t$. So, we have:

Theorem 3.1 *If $k = 2, 3$, or 4 , then*

$$\nu_p(b_m(y_m^k)) = \max\{r - \lfloor \frac{2(k-1)}{p-1} \rfloor + \nu_p(r) : \lfloor \frac{2k}{p-1} \rfloor \leq r \leq \lfloor \frac{m}{p-1} \rfloor\}$$

for each $p \geq 2$. Further, this formula is true if $p = 2, 3$ and $k = 5, \dots, t$ or if $k = 5$, or 6 and $p > 5$.

REMARK. From the above discussion, it is reasonable to conjecture that :
If p is any prime number and $k = 5, \dots, t$, then

$$\nu_p(b_m(y_m^k)) = \max\{r - \lfloor \frac{2(k-1)}{p-1} \rfloor + \nu_p(r) : \lfloor \frac{2k}{p-1} \rfloor \leq r \leq \lfloor \frac{m}{p-1} \rfloor\}.$$

Now, we compute the J -order of any complex line bundle over $\mathbb{C}P^m$.

EXAMPLE 2. Let $n \in \mathbb{N}$. Then the J -order of $\xi_m(\mathbb{C})^n$ is $b_m(r\xi_m(\mathbb{C})^n - 2)$.
By Theorem 2.5, $\nu_p(b_m(r\xi_m(\mathbb{C}) - 2)) = \nu_p(e_t(\theta_p(r\xi_m(\mathbb{C})^n - 2)))$.

$$r\xi_m(\mathbb{C})^n - 2 = r\psi^n(\xi_m(\mathbb{C}) - 1) = \psi^n(r\xi_m(\mathbb{C}) - 2) = \psi^n(y_m).$$

So, by Corollary 2.3,

$$\theta_p(r\xi_m(\mathbb{C})^n - 2) = \theta_p(\psi^n(y_m)) = \psi^n(\theta_p(y_m)).$$

Let $n = p_1^{r_1} \dots p_s^{r_s} p^d$ where $r_i > 0$ for $i = 1, \dots, s$, and $d \geq 0$. Let $p = 2q + 1$ be any odd prime number. By, Lemma 5.4 [7],

$$\theta_p(y_m) = 1 + \sum_{j=1}^{q-1} m_j y_m^j + \frac{1}{p} y_m^q$$

where $m_j \in \mathbb{Z}$ with $\nu_p(m_j) = 0$ for $j = 1, \dots, q-1$. Hence,

$$\psi^n(\theta_p(y_m)) = 1 + \sum_{j=1}^{q-1} m_j \psi^n(y_m)^j + \frac{1}{p} \psi^n(y_m)^q.$$

Using Lemma 3.6 of [7], we easily obtain that

$$\psi^{p^d}(y_m) = \sum_{j=1}^{p^d-1} n_j y_m^j + y_m^{p^d}$$

with $\nu_p(n_j) > 0$ for $j = 1, \dots, p^d - 1$. Now, using the fact that Adams operations are ring homomorphisms with $\psi^{l_1} \circ \psi^{l_2} = \psi^{l_1 l_2}$ for each $l_1, l_2 \in \mathbb{Z}$, we get $\psi^n(y_m) = \sum_{j=1}^n a_j y_m^j$ where $a_j \in \mathbb{Z}$ such that $\nu_p(a_j) = 0$ if j is a multiple of p^d , and $\nu_p(a_j) \geq 1$ for all other values of j . Hence,

$$\psi^n(\theta_p(y_m)) = 1 + \sum_{j=1}^{nq} b_j y_m^j$$

such that $\nu_p(b_j) = -1$ if j is a multiple of $p^d q$, $\nu_p(b_j) \geq 1$ for $j < p^d q$, and $\nu_p(b_j) \geq 0$ for all other values of j . Hence, $\theta_p(\psi^n(y_m))$ is a strict D-series at p of type (α, β) where

$$\alpha_{p'} = \begin{cases} p^d q & p' = p \\ \infty & p' \neq p \end{cases}, \quad \beta_{p'} = \begin{cases} 1 & p' = p \\ \infty & p' \neq p. \end{cases}$$

Hence,

$$\nu_p(b_m(r\xi_m(\mathbb{C})^n - 2)) = \max\{r + \nu_p(r) : 0 \leq r \leq [\frac{m}{p^{\nu_p(n)}(p-1)}]\}.$$

If $p = 2$, then

$$\theta_p(y_m) = (1 + \frac{1}{4}y_m)^{\frac{1}{2}}.$$

So,

$$\psi^n(\theta_p(y_m)) = (1 + \frac{1}{4}\psi^n(y_m))^{\frac{1}{2}}.$$

Let $n = p_1^{r_1} \dots p_s^{r_s} 2^d$. Then $1 + \frac{1}{4}\psi^n(y_m) = \sum_{j=1}^n c_j y_m^j$ with $\nu_p(c_j) = -2$ if j is a multiple of 2^d , $\nu_p(c_j) \geq 0$ for $j < 2^d$, and $\nu_p(c_j) \geq -1$ for all other values of j . Hence, $1 + \frac{1}{4}\psi^n(y_m)$ is a strict D-series at 2 of type (α, β) where

$$\alpha_{p'} = \begin{cases} 2^d & p' = 2 \\ \infty & p' \neq 2 \end{cases}, \quad \beta_{p'} = \begin{cases} 2 & p' = 2 \\ \infty & p' \neq 2. \end{cases}$$

So,

$$\nu_2(e_t(1 + \frac{1}{4}\psi^n(y_m))) = \max\{2r + \nu_2(r) : 0 \leq r \leq [\frac{t}{2^d}]\}.$$

Hence,

$$\begin{aligned} \nu_2(b_m(r\xi_m(\mathbb{C})^n - 2)) &= \nu_2(e_t((1 + \frac{1}{4}\psi^n(y_m))^{\frac{1}{2}})) \\ &= \max\{2r + \nu_2(2r) : 0 \leq r \leq [\frac{t}{2^d}]\} = \max\{r + \nu_2(r) : 0 \leq r \leq [\frac{m}{2^d}]\}. \end{aligned}$$

So, we have:

Theorem 3.2 *Let $n \in \mathbb{N}$ and p be any prime number then*

$$\nu_p(b_m(r\xi_m(\mathbb{C})^n - 2)) = \max\{r + \nu_p(r) : 0 \leq r \leq [\frac{m}{p^{\nu_p(n)}(p-1)}]\}.$$

REMARK. A similar proof of Theorem 3.2 when n is a power of a prime p has been obtained independently by Dibağ [5].

References

- [1] Adams, J.F.: On the groups $J(X)$, I, II, III, *Topology* **2** (1963), 181–196, **3** (1965), 137–171, 193–222.
- [2] Adams, J.F. and Walker, G.: On complex Stiefel manifold, *Proc. Camb. Phil. Soc.* **56** (1965), 342–353.
- [3] Atiyah, M.F.: Thom complexes, *Proc. London Math. Soc.* (3) **11** (1961), 291–310.
- [4] Dibağ, I.: Integrality of rational D -series, *J. Algebra* (2) **164** (1994), 468–480.
- [5] Dibağ, I.: Proof of Lam’s results, preprint.
- [6] Lam, K.Y.: Fibre homotopy trivial bundle over complex projective spaces, *Proc. Amer. Math. Soc.* **33** (1972), 211–212.
- [7] Muhammad, N., Piccinini, R., and Suter, U.: Some applications of topological K-theory, *North-Holland Math. Studies*, No.45, North Holland, Amsterdam, 1980.
- [8] Obiedat, M.: A note on the localization of J -groups, to appear in *Hiroshima Math. J.*, **2** (1999).
- [9] Quillen, D.: The Adams conjecture, *Topology* **10** (1970), 67–80.

University of California
Department of Mathematics
Berkeley, CA 94720, USA
e-mail : obiedat@math.berkeley.edu